



Volume 3 · Issue 1 · March / April 2013

Geoinformatica - An International Journal (GIJ)

ISSN : 2180-1231

Number of issues per year: 6

CSC PUBLISHERS

<http://www.cscjournals.org>

Geoinformatica – An International Journal (GIJ)

VOLUME 3, ISSUE 1, 2013

**EDITED BY
DR. NABEEL TAHIR**

ISSN (Online): 2180-124X

Geoinformatica – An International Journal (GIJ) is published both in traditional paper form and in Internet. This journal is published at the website <http://www.cscjournals.org>, maintained by Computer Science Journals (CSC Journals), Malaysia.

GIJ Journal is a part of CSC Publishers

Computer Science Journals

<http://www.cscjournals.org>

Geoinformatica – An International Journal (GIJ)

Book: Volume 3, Issue 1, March / April 2013

Publishing Date: 30-04-2013

ISSN (Online): 2180-124X

This work is subjected to copyright. All rights are reserved whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, re-use of illustrations, recitation, broadcasting, reproduction on microfilms or in any other way, and storage in data banks. Duplication of this publication of parts thereof is permitted only under the provision of the copyright law 1965, in its current version, and permission of use must always be obtained from CSC Publishers.

GIJ Journal is a part of CSC Publishers

<http://www.cscjournals.org>

© GIJ Journal

Published in Malaysia

Typesetting: Camera-ready by author, data conversion by CSC Publishing Services – CSC Journals, Malaysia

CSC Publishers, 2013

EDITORIAL PREFACE

Geoinformatica – An International Journal (GIJ) is an effective medium for interchange of high quality theoretical and applied research in Geoinformatica domain from theoretical research to application development. This is the *First* Issue of Volume *Three* of GIJ. The Journal is published bi-monthly, with papers being peer reviewed to high international standards. GIJ emphasizes on efficient and effective geomatic sciences, and provides a central for a deeper understanding in the discipline by encouraging the quantitative comparison and performance evaluation of the emerging components of Geoinformatica. Some of the important topics are spatial ontologies, computational geometry and visualization for geographic information systems, geostatistics and spatial statistics, spatial analysis, interoperability, and innovative applications of geotechnologies etc.

The initial efforts helped to shape the editorial policy and to sharpen the focus of the journal. Started with Volume 3, 2013, GIJ appear with more focused issues. Besides normal publications, GIJ intend to organized special issues on more focused topics. Each special issue will have a designated editor (editors) – either member of the editorial board or another recognized specialist in the respective field.

GIJ give an opportunity to scientists, researchers, and vendors from different disciplines of Geoinformatica to share the ideas, identify problems, investigate relevant issues, share common interests, explore new approaches, and initiate possible collaborative research and system development. This journal is helpful for the researchers and R&D engineers, scientists all those persons who are involve in Geoinformatics in any shape.

Highly professional scholars give their efforts, valuable time, expertise and motivation to GIJ as Editorial board members. All submissions are evaluated by the International Editorial Board. The International Editorial Board ensures that significant developments in geotechnologies from around the world are reflected in the GIJ publications.

GIJ editors understand that how much it is important for authors and researchers to have their work published with a minimum delay after submission of their papers. They also strongly believe that the direct communication between the editors and authors are important for the welfare, quality and wellbeing of the Journal and its readers. Therefore, all activities from paper submission to paper publication are controlled through electronic systems that include electronic submission, editorial panel and review system that ensures rapid decision with least delays in the publication processes.

To build its international reputation, we are disseminating the publication information through Google Books, Google Scholar, Directory of Open Access Journals (DOAJ), Open J Gate, ScientificCommons, Docstoc and many more. Our International Editors are working on establishing ISI listing and a good impact factor for GIJ. We would like to remind you that the success of our journal depends directly on the number of quality articles submitted for review. Accordingly, we would like to request your participation by submitting quality manuscripts for review and encouraging your colleagues to submit quality manuscripts for review. One of the great benefits we can provide to our prospective authors is the mentoring nature of our review process. GIJ provides authors with high quality, helpful reviews that are shaped to assist authors in improving their manuscripts.

Editorial Board Members

Geoinformatica – An International Journal (GIJ)

EDITORIAL BOARD

EDITORIAL BOARD MEMBERS (EBMs)

Dr. Lim Hwee San
University Sains Malaysia
Malaysia

TABLE OF CONTENTS

Volume 3, Issue 1, March / April 2013

Pages

- | | |
|--------|--|
| 1 - 11 | Multi-Scale Urban Analysis Using Remote Sensing and GIS
<i>P. Kalyani, P. Govindarajulu</i> |
|--------|--|

A multi-scale Urban Analysis Using Remote Sensing and GIS

P.Kalyani

*Dept. of Computer Applications,
Priyadarshini College of Engineering
Nellore, AndhraPradesh,India.*

kalyani.purini@gmail.com

P.Govindarajulu

*Dept. of Computer Science,
S.V. University
Tirupati, AndhraPradesh,India.*

pgovindarajulu@yahoo.com

Abstract

Urban planning was very much a design and engineering exercise with the state as a single stake holder. Mega cities with millions of population, has undergone a series of physical as well as socio-economic changes over the last 60 years. In India, Hyderabad experienced a high rate of urbanization facing structural, environmental, social and economic problems. To provide a holistic perspective on the urban characteristics, an interdisciplinary research approach is used. GIS-Geographic Information System and Remote Sensing provide the advance techniques and methods for studying urban land development and assist urban planning.

Keywords: GIS, Remote Sensing, Multi-Scale Urban Analysis, Land Use Change, Urban Growth, Urban Planning.

1. INTRODUCTION

Mega cities the largest category of urban agglomeration, attract considerable attention because of their population size, economic, socio-cultural, environmental and political influence and geographical complexity. Until 1975 there were just three mega cities in the world. The number of so-called mega cities increased in the period from 1975 until today from 4 to 22, mostly in less developed regions [11]. The number of cities increased to 27 mega cities until 2015 [8]. The population development of the world is expected to increase continuously from 6.7 billion to 9.3 billion in 2050. But a heavy increase of mega cities creates a serious problem in India. The population of India (today 1.2 billion) has grown two and half times, but the urban population has grown nearly five times. The number of Indian mega cities will double from the current three (Mumbai, Delhi and Kolkata) to six by the year 2021, (new additions will be Bangalore, Chennai and Hyderabad), when India will have the largest concentration of mega cities in the world [4].

Then the number of six mega cities (Mumbai, Delhi, Kolkata, Bangalore, Chennai and Hyderabad) is increased by twelve by the year 2015 (new Ahmadabad, Pune, Surat, Kanpur, Jaipur and Lucknow) [17]. With the rapid increase of urban growth, makes us to face lot of socio-economic, environmental and political problems. This phenomenon will necessitate advanced methodologies such as space technologies, which help city planners, economists, environmentalists, ecologists and resource managers solve the problems which accompany such growth [9]. Urban planners need information about the rate of growth, pattern and extent of sprawl to provide the basic amenities such as water, sanitation and electricity etc. Since planners currently lack such information, most of the sprawl areas lack basic infrastructure facilities.

On the last decade, earth observation sensors developed to a stage where global maps have been made possible on low resolution (LR) from 250m to 2 Km [14]. Examples are global urban extent maps based on e.g., DMSP-OLS night-time lights imagery [6], MODIS data [3][15]. A list,

analysis and comparison of the various available global data sets is presented and discussed by Potere and Schneider [13]. However, most of them are provided for a single time step, and the coarse geometric resolution is a clear restriction tracing the small-scale urban outlines, extents and patterns. Even though higher resolution sensors systems are available e.g., Land sat, spot, Rapid Eye, IRS, IKONOS, Quick Bird, World View-I and II. The provision of a global coverage or at least of a large amount of cities – is not an easy task. Limitations such as cloud coverage, on board storage capacity, sensor utilization and sharing of the same source with other EO projects cause a several years lasting acquisition period. Furthermore, data costs and processing effort are significant. Thus, a global coverage at the scale covered by the medium (MR: here defined as on 10m to 100m) and high resolution (HR: 1M-10m) to very high resolution (VHR: <1m). EO sensors are inexistence.

Research studies on long term monitoring of the spatial effects of the urbanization are mostly based on MR (Medium Resolution) data from sensors such as Land sat or spot, having lower geometric resolution and thus allow for fewer thematic details. Different studies have also shown that radar imagery is an excellent basis for classifying, monitoring and analyzing urban conglomerations and their development overtime especially in cases of large area mapping [5]. Using of MSS (Multi Spectral Scanner) data, ETM (Enhanced Thematic Mapped data) and Terra SAR-X Strip map data is used for monitoring urbanization in mega cities from space for analysis of 22 to 27 mega cities and their number is constantly increasing [8]. Temporal and spatial urban sprawl, re-densification and urban development in the tremendously growing six mega cities to 12 mega cities in India, and became the largest urban agglomerations [19].

In India, by using Quick bird data of VHR (Very High Resolution $\leq 1\text{M}$ i.e., 0.61M) with a sub-meter geo-metric resolution is applied for the multi-scale urban analysis of the Hyderabad metropolitan area of deriving parameters such as houses, streets, shadows, vegetation, bare soil etc., [17]. For the analysis of the urban patterns, first we have to classify the obtained data. The classification of the various land-sat scenes is based on an object-oriented classification procedure [16]. The first step is a multi-resolution segmentation. The second step is a hierarchical thematic classification procedure allowing mapping four different thematic classes, namely 'water', 'vegetation', 'undeveloped land' and 'urban' [2].

In this paper focused on the multi-scale approach with remote sensing, to support urban management with area-wide and up-to-date datasets in Hyderabad.

The main objective of this paper is:

- Landsat data enable to monitor the entire urban area and its extension as overview level.
- A time series allows for urban change detection to study urban growth and its characteristics overtime.
- At the far end of the multi-level approach Quick bird data provide highly detailed information on the structural characteristics of the urban morphology.

Further on, it will be illustrated how remote sensing can contribute basic information for improving technical supply and disposal infrastructure, e.g. for water, transportation, energy, sewage water, waste, traffic etc.

2. STUDY AREA

India is a large country, whose population is concentrated predominantly along the Mediterranean climate, is close proximity to its major cities Delhi, Kolkata, Mumbai, Chennai, Bangalore, Hyderabad figure(1). The study focuses on Hyderabad city, is a sprawling metropolis with a population of 6.8 million and growth rate of 2.5 percent per year. The population of the city can be reached to 10 to 13 million in 2020 turning it into one of the megacities of tomorrow [12].

“Megacities are undergoing new dynamics and, as a consequence are facing new spatial and organizational challenges” [1]. The performance in megacities is seen as a key factor regarding global sustainable development .First targets for future development are set in Hyderabad’, master plan and in the “City Development Strategy” of 2003 [12]. The city has faced many problems such as

- Transport system
- Water supply
- Pollution of air, water, sound etc
- Inner city infrastructure etc




FIGURE 1: Location of Hyderabad in India.

3. DATA AND METHODS

A. Data

It started with the Multispectral-scanner(MSS) featuring a geometric resolution of 59 meters and a spectral resolution of four bands(green, red, two near infrared bands).Since 1982 the thematic Mapper (TM) has operated with 30 m geometric resolution and seven spectral bands. Since 1999 the Enhanced Thematic Mapper (ETM) has operated with an additional panchromatic band and 15m geometric resolution. Since 2002 Ikonos data of with 1m geometric resolution and since 2005 Quick bird data with 0.61m geometric resolution for finding the illegal constructions in the inner city [17].With its field of view of 185 KM the satellite is able to survey the large metropolitan areas of the study sites-thus covering in dependence of their spatial position entire areas and no cloud coverage.

The Landsat images 1997 and 2007 [22] [21] are collected from NRSC; Hyderabad is used for to detect the urban growth. The other Quick bird data of the year 2010 is also collected from NRSC; Hyderabad [20] is used for high resolution analysis.

The level of description with Land sat features is not flooded with microscopic detail, but re-gives nevertheless the specific features of the urban system. For this purpose, the requirements for the differentiation of classes are limited to the classification of built-up and non-built-up areas.

B. Methodologies for Land Use / Land Cover Classification

Landsat data of Hyderabad for different years were used for the detection of recent changes of the urban extension. The data was enhanced before classification using histogram equalization in ERDAS Image 9.2 to improve the image quality and to achieve better classification accuracy. A land cover classification extracting the classes' built-up areas, non-built-up areas, vegetation and water was performed separately on both images. The main goal is to identify the urban built-up areas to measure the changes of the urban extension over the time interval. For that purpose the classification methodology is based on an object-oriented hierarchical approach. The object-oriented methodology was used to combine spectral features with shape, neighborhood and texture features [17].

Post classification comparison was found to be the most accurate procedure and presented the advantage of indicating the nature of the changes [10]. An automated object-oriented for an extraction of detached houses, of main street infrastructure, of vegetation areas, of bare soil and of water areas. The methodology presented by [17] was applied to the Quick Bird data of Hyderabad.

A comparative analysis of land cover classification analysis of land cover classifications for times t_1 and t_2 performed independently was therefore implemented to monitor and analyze the land cover changes in the metropolitan area of Hyderabad. Pixel wise change detection was implemented in the flow chart figure (2) checking the land cover classes individually.




FIGURE 2: Flow Chart of Methodology for Land Use/Land Cover and Change Detection.

4. RESULTS, CHANGE DETECTION AND DISCUSSIONS

For the mega cities of the world featuring four individual urban foot print classification at four time steps in the mid-1970's, around 1990, 2000 and 2010. The change detection allows identifying, localizing and quantifying the pattern and dimension of urban sprawl overtime. Since 1975 the population development and thus, the rate of urbanization in the mega cities outside the high developed countries were enormous [8]. At a first glance, a number of trends are immediately clear in terms of spatial dynamics of urbanization over the time period observed is the highest in developing countries. Beyond this, the spatial extents in developed countries are noticeably larger with respect to the absolute population. The first and the most natural analysis is the measurement of spatial urban expansion over time. The quantification of urban growth for the metropolitan areas of the mega cities are calculates as relative growth.

The urban analysis aims at a holistic understanding of the dynamics of urban growth process. The different perspectives on the city of Hyderabad are caused through the different data types. In contrary the capabilities of the Quick Bird data with its high geometry resolution allow highly detailed analysis of the urban structure and the neighborhoods [17]. The different remote sensing data contribute to a multi-level perspective within an interdisciplinary GIS database to support the information basis for sustainable urban development.

A. Remote sensing

Here we used the images of Hyderabad of the years 1997 and 2007 for the detection of recent changes of the urban extension. The contribution of remote sensing to the planning of sustainable urban development is two-fold. On the one end hand a spatial overview on the extension of the city and its structure change over time is presented. On the other hand a highly detailed analysis of the urban structure.




FIGURE 3: Landsat Imagery of Hyderabad, 2007.

The main goal is to identify the urban built-up areas to measure the changes of the urban extension over the time interval. A comparative analysis of land cover classification for times t1 and t2 performed independently was therefore implemented to monitor and analyze the land use patterns in the metropolitan area of Hyderabad. Pixel wise change detection was implemented checking the land cover classes individually of the two available years.

In figure (4), the spatial expansion of the urban built-up areas of Hyderabad between 1997 and 2007. In particular, the urban sprawl in the periphery and densification process in Hyderabad. New developments arise on the outskirts of Hyderabad. Nevertheless, focal points emerge in the northeast as well as in the southeast shows in the results.




FIGURE 4: Urban Growth in Hyderabad Between 1997 and 2007.

In 1997, the urbanized area was 130km². By the change detection, the urbanized areas were increased of about 2.5 times as large as in 1997 was measured. In the olden days only low resolution data is available. For a large scale analysis of urban growth in the metropolitan area of Hyderabad high resolution satellite image i.e. QuickBird data, Ikonos data were came up. That satellite image can support a detailed analysis of the small-scale urban morphology. In addition to this example a satellite data of another time period i.e. 2010, an automated object-oriented classification approach was implemented for an extraction of detached houses, of main street infrastructure, of vegetation areas of bare soil and of water areas. The methodology presented by [17] was applied to the QuickBird data of Hyderabad.

The methodical framework was inherited, with adjustments on the spectral peculiarity of the particular sense. For validation of the classification results an accuracy assessment has been carried out. The accuracy shows an average of more than 80% of houses have been classified correctly. This analysis of the current status of the built-up environment provides useful spatial information for a sustainable urban management. In future enables the calculation of the built-up density and provides necessary information for the analysis of the location and the carrying capacity of the street infrastructure as well as the analysis of accessibilities. The main infrastructure was extracted for an assessment of the street network performance against built-up densities. Highly dense built-up areas alternate with less dense areas, or even open spaces. The urban land covers classification of the heterogeneous urban structure in Hyderabad shown below.



FIGURE 5: QuickBird Imagery of Hyderabad, 2010.

Noticeable are the different urban morphologies which are a result of the various building sizes, shapes or alignments. The shape features enable a basic differentiation between residential and commercial usage.

Remote sensing is a feasible instrument for efficient and cost-effective analysis of the urban land cover overtime. Urban structures reflect the distribution of the population, the level of prosperity, and different units as well as infrastructure. These results from remote sensing are indispensable up-to-date spatial information to support substantial urban planning and decision making.

B. Interdisciplinary Integration

The infrastructure for housing, vegetation, water bodies etc is not sufficient, but we want spatial information for future planning. This infrastructure is not directly be measured from space, we also collected data by field surveys showing the urban structure, social structure, mobility patterns, energy and water flows. These data will be combined with remote sensing in order to expand this local information to a wider area. The results will be organized in a common GIS database. A detailed analysis of demand is a prerequisite for technical infrastructure planning of both supply and disposal services, e.g., of water, energy, sewage water and waste. The development often lack a cross cutting analyses also assessing the impact on different sectors. A common database is a first step to provide necessary information. Procedures for cross cutting analyses and evaluation have to be developed and introduced. The first cross analyses with existing data coming from the master plan in selected neighborhood show already the frictions. But available data from the city administration are not up to date and new illegal and informal constructions can be found all over the city.

Since large as well as small and medium sized man made constructions which are numerous in the city of Hyderabad are important users of technical infrastructure, a detailed knowledge of their demand and spatial distribution would allow an infrastructure planning that a better adapted to the real needs. It can also serve as basic information for improving technical equipment concerning traffic in the roads, water use, waste, energy etc and thus achieving resource savings and emission reductions. Therefore, a GIS based inventory and including information on their demand of supply and disposal services is being developed for a part of the city of Hyderabad. Remote sensing provides a first indication of the position of urban growths arise on the outskirts of Hyderabad.




FIGURE 6: Analysis of services in Telecom colony based on remote sensing data supported by site visits.

For an inventory of different built-up areas clearly, the remote sensed data is supplemented by information provided by local partners are collected in surveys and site visits which allows in the end an assessment of supply and disposal needs. A possible example of the application of remote sensing combined with site visits to analyze the situation in the “Telecom colony” neighborhood, at Hi-tech city, e.g. education, worship, commercial etc. The land use data (green and orange) in figure (6) is derived from remote sensed data and represents the basis for a mapping sensed data and represents the basis for a mapping of the services. Hence, the interdisciplinary approach enables a more holistic view on the urban functions and structures. Beside the basic data about the situation today also monitoring is a big issue. Therefore one of the basic questions is what kind of data is really necessary and can be kept up to date with a reasonable effort. Taking into account the rapid changes, methods like remote sensing are expected to bring a substantial contribution.

5. CONCLUSION

To fight with the problems faced by the rapid urban growth, various scientific perspectives are necessary for a holistic assessment of urban situations. The major problems associated with the urban centers in India are that of unplanned expansion, changing land use / land cover areas [4].

For this, remote sensing imagery, with its repetitive and synoptic viewing capabilities, together with GIS, is important tools to map areas and monitor the changes in the urban growth. High-resolution satellite data is applied for the multi-scale urban analysis of the Hyderabad metropolitan area of deriving parameters such as houses, streets, shadows, vegetation, bare soil etc., [17]. This paper focused on the multi-scale approach with remote sensing, to support urban management with area-wide and up-to-date datasets. The Land sat analysis enabled the monitoring of the urban growth overtime to understand the dynamics and characteristics of the metropolitan area of Hyderabad. This includes also information about the direction of urban growth.

Thus, remote sensing and GIS provided to be a very useful basis for a more detailed analysis of the spatial distribution, an emerging megacity which is prerequisite for a reasonable planning of technical infrastructure. The officials of various government departments should be given through exposure and training of remote sensing and GIS for its application implementation in the urban

management plans. This multilayer spatial information allows analyzing and anticipating developments to support future planning strategies.

6. REFERENCES

- 1 A. G. Aguilar, P. M. Ward: Globalization, regional development, and mega-city expansion in Latin America, loc. cit., p. 4. Hyderabad Urban Development Authority (HUDA), January 2003, "Hyderabad 2020, Plan for sustainable development", Draft Master Plan for Hyderabad Metropolitan Area.
- 2 Abelen, S. (2010). Development of a user interface for optimizing urban area classification from Landsat data. Unpublished Master thesis, Technical University Munich;p. 111.
- 3 Bartholome, E., & Belward, A. S. (2005). GLC2000: A new approach to global land covers mapping from Earth observation data. *International Journal of Remote Sensing*, 26. (pp. 1959–1977).
- 4 Chakrabati, P. G. D. (2001). Urban crisis in India: new initiatives for sustainable cities. *Development in Practice*, 11(2–3), 260–272.
- 5 Dell'Acqua, F. (2009). The role of SAR sensors. In P. Gamba, & M. Herold (Eds.), *Global mapping of human settlements: Experiences, data sets, and prospects* (pp. 209–319). Boca Raton: Taylor and Francis
- 6 Elvidge, C., Imhoff, M. L., Baugh, K. E., Hobson, V. R., Nelson, I., Safran, J., et al. (2001). Nighttime lights of the world: 1994–95. *ISPRS Journal of Photogrammetry and Remote Sensing*, 56, 81–99.
- 7 Esch, T., Thiel, M., Schenk, A., Roth, A., Mehl, H., & Dech, S. (2010, Feb.). Delineation of urban footprints from TerraSAR-X data by analyzing speckle characteristics and intensity information. *IEEE Transactions on Geoscience and Remote Sensing*, 48(2), 905–916.
- 8 H.Taubenbock,T.Esch,A.Felbier,M.Wiesner,A.Roth,,S.Dech(2012):Monitoring Urbanization inMegacities from space.remote sensing of Environment,117,162-176.
- 9 Maktav, D., & Erbek, F. S. (2005). Analyse of urban growth using multitemporal satellite data in Istanbul, Turkey. *International Journal of Remote Sensing*, 26(4).*International Journal of Remote Sensing*, 26(4).
- 10 MAS, J.-F. (1999): Monitoring land-cover changes: a comparison of change detection techniques. *International Journal of Remote Sensing*, vol. 20, No. 1, pp. 139-152.
- 11 Münchner Rück (2005). Megastädte – Megarisiken. Trends und Herausforderungen für Versicherung und Risikomanagement. www.munichre.com/publications/302-04270_de.pdf.
- 12 Municipal Corporation of Hyderabad (MCH), August 2003, "City Development Strategy Hyderabad: Strategic action plan and city assistance programme", www.cdshyderabad.org.in.

- 13 Potere, D., & Schneider, A. (2009). Comparison of global urban maps. In P. Gamba, & M. Herold (Eds.), *Global mapping of human settlements: Experiences, data sets, and prospects* (pp. 269–308). Taylor & Francis Group.
- 14 Potere, D., Schneider, A., Angel, S., & Civco, D. L. (2009). Mapping urban areas on a global scale: Which of the eight maps now available is more accurate? *International Journal of Remote Sensing*, 30(24), 6531–6558.
- 15 Schneider, A., Friedl, M., & Woodcock, C. (2005). Mapping urban areas by fusing multiplesources of coarse resolution remotely sensed data: Global results. Proc. 5th int.symp. Of remote sensing of urban areas (Tempe, AZ, March 2005).
- 16 Taubenböck, H. (2008). *Vulnerabilitätsabschätzung der Megacity Istanbul mit Methoden der Fernerkundung*. Ph.D. thesis. University of Würzburg; p. 178. ISBN-10: 3639083180.
- 17 Taubenböck, H., Pengler, I., Schwaiger, B., Cypra, S., Hiete, M., & Roth, A. (2007). A multi-scale urban analysis of the Hyderabad Metropolitan area using remote sensing and GIS (p. 6). Urban remote sensing joint event, Paris, France.
- 18 Taubenböck, H., Roth, A., 2007, A transferable and stable object oriented classification approach in various urban areas and various high resolution sensors. Urban Remote Sensing Joint Event, Paris, France.
- 19 Taubenböck, H., Wegmann, M., Roth, A., Mehl, H., & Dech, S. (2009). Urbanization in India — Spatiotemporal analysis using remote sensing data. *Computers, Environment and Urban Systems*, 33, 179–188.
- 20 United Nations, 1997, *World Urbanization Prospects, The 1997 Revision*, New York.
- 21 United Nations, 2007, *World Urbanization Prospects, The 2007 Revision*, New York.
- 22 United Nations, 2010, *World Urbanization Prospects, The 2010 Revision*, New York.

INSTRUCTIONS TO CONTRIBUTORS

Geoinformatica – An International Journal (GIJ) aims at publishing scientific and technical developments in the diverse field of Geoinformatics. GIJ covers all aspects and information on scientific and technical advances in the geomatic sciences. The journal is providing a platform for exploring research, development and innovative applications in geographic information science and related areas. GIJ provides a privileged view of what is currently happening in the field of geoinformatics as well as a preview of what could be the hottest developments and research topics in the near future. Additionally, it includes recent research results on spatial databases, spatial ontologies, computational geometry and visualization for geographic information systems, geostatistics and spatial statistics, spatial analysis, interoperability, and innovative applications of geotechnologies.

To build its International reputation, we are disseminating the publication information through Google Books, Google Scholar, Directory of Open Access Journals (DOAJ), Open J Gate, ScientificCommons, Docstoc and many more. Our International Editors are working on establishing ISI listing and a good impact factor for GIJ.

The initial efforts helped to shape the editorial policy and to sharpen the focus of the journal. Started with Volume 3, 2013, GIJ appear with more focused issues. Besides normal publications, GIJ intend to organized special issues on more focused topics. Each special issue will have a designated editor (editors) – either member of the editorial board or another recognized specialist in the respective field.

We are open to contributions, proposals for any topic as well as for editors and reviewers. We understand that it is through the effort of volunteers that CSC Journals continues to grow and flourish.

LIST OF TOPICS

The realm of Geoinformatica – An International Journal (GIJ) extends, but not limited, to the following:

- Applied Geography
- Close Range and Videometric Photogrammetry
- Computational Geometry and Visualization
- Distributed GIS/GIS and the Internet
- Geodata: Capture, Sources and Standards
- Geographic Information Science
- Geospatial Web
- Geostatistics
- Guidance Systems
- Land and Geographic Information Systems
- Location-Based Services
- Mobile Maps
- Sensor Networks
- Spatial Cognition
- Spatial Ontologies and Interoperability
- Surface Modeling
- Digital Mapping
- Geo Tags
- Geographic Data
- Geographic Information
- Geoinformatics
- Geospatial Applications
- Geospatial Databases
- Geospatial Processing
- Global Positioning System
- Integrated Geodesy
- Web Services
- Map Services
- Remote Sensing
- Sensor Web
- Spatial Data Analysis
- Spatial Databases

CALL FOR PAPERS

Volume: 3 - Issue: 2

i. Paper Submission: April 30, 2013

ii. Author Notification: May 31, 2013

iii. Issue Publication: June 2013

CONTACT INFORMATION

Computer Science Journals Sdn Bhd

B-5-8 Plaza Mont Kiara, Mont Kiara

50480, Kuala Lumpur, MALAYSIA

Phone: 006 03 6204 5627

Fax: 006 03 6204 5628

Email: cscpress@cscjournals.org

CSC PUBLISHERS © 2013
COMPUTER SCIENCE JOURNALS SDN BHD
B-5-8 PLAZA MONT KIARA
MONT KIARA
50480, KUALA LUMPUR
MALAYSIA

PHONE: 006 03 6204 5627
FAX: 006 03 6204 5628
EMAIL: cscpress@cscjournals.org